

DECLARATION

I, Makoto AIHARA, Patent Attorney, of SIKs & Co., 8th Floor, Kyobashi-Nisshoku Bldg., 8-7, Kyobashi 1-chome, Chuo-ku, Tokyo 104-0031 JAPAN hereby declare that I am the translator of the certified official copy of the documents in respect of an application for a patent filed in Japan on January 30, 2002 under Patent Application No. 021159/2002 and that the following is a true and correct translation to the best of my knowledge and belief.

Dated: November 20, 2006


Makoto AIIARA

[Title]	Application for Patent
[Reference Number]	A21046A
[Filing Date]	January 30, 2002
[Address]	To the Commissioner of the Patent Office
[Inventor]	
[Domicile or Residence]	c/o RIKEN, 2-1, HiroSawa, Wako-shi, Saitama
[Name]	Shou TAKASHIMA
[Inventor]	
[Domicile or Residence]	c/o RIKEN, 2-1, HiroSawa, Wako-shi, Saitama
[Name]	Masafumi TSUJIMOTO
[Inventor]	
[Domicile or Residence]	2-26, Muromachi, Hadano-shi, Kanagawa
[Name]	Shuichi TSUJI
[Applicant for Patent]	
[Identification Number]	000006792
[Name]	RIKEN
[Administrator]	
[Identification Number]	100092635
[Patent Attorney]	
[Name]	Hisao SHIOZAWA
[Subadministrator]	
[Identification Number]	100096219
[Patent Attorney]	
[Name]	Masazumi IMAMURA
[Subadministrator]	
[Identification Number]	100095843
[Patent Attorney]	
[Name]	Junji KAMATA
[Fee]	
[Number of register of payment]	007663
[Amount]	21000
[List of Attached Documents]	
[Document's Name]	Specification 1
[Document's Name]	Drawing 1
[Document's Name]	Abstract 1
[Number of Comprehensive Power of Attorney]	9607613
[Proof]	Required

[Name of Document] SPECIFICATION
[Title of Invention] GLYCOSYLATING ENZYME
[Claims]

[Claim 1] O-glycan α 2,8-sialyltransferase, which is characterized in that it has the following substrate specificity and substrate selectivity.

Substrate specificity: the substrates of the enzyme are glycoconjugates having a Sia α 2,3(6)Gal structure (wherein Sia represents sialic acid and Gal represents galactose) at the terminus thereof.

Substrate selectivity: the enzyme incorporates sialic acids into O-glycans more preferentially than into glycolipids or N-glycans.

[Claim 2] O-glycan α 2,8-sialyltransferase having either one of the following amino acid sequences:

- (1) an amino acid sequence shown in SEQ ID NO: 1; or
- (2) an amino acid sequence comprising a deletion, substitution, and/or addition of one or several amino acids with respect to the amino acid sequence shown in SEQ ID NO: 1, and having O-glycan α 2,8-sialyltransferase activity.

[Claim 3] O-glycan α 2,8-sialyltransferase gene encoding the amino acid sequence of the O-glycan α 2,8-sialyltransferase according to claim 2.

[Claim 4] The O-glycan α 2,8-sialyltransferase gene according to claim 3 which has any one of the following nucleotide sequences:

- (1) a nucleotide sequence corresponding to a portion between nucleotide 77 and nucleotide 1270 of a nucleotide sequence shown in SEQ ID NO: 2; or
- (2) a nucleotide sequence comprising a deletion, substitution, and/or addition of one or several nucleotides with respect to the nucleotide sequence corresponding to a portion between nucleotide 77 and nucleotide 1270 of the nucleotide sequence shown in SEQ ID NO: 2, and encoding a protein having O-glycan α 2,8-sialyltransferase activity;

[Claim 5] A recombinant vector comprising the O-glycan α 2,8-sialyltransferase gene according to claim 3 or 4.

[Claim 6] The recombinant vector according to claim 5 which is an expression vector.

[Claim 7] A transformant transformed with the recombinant vector according to claim 5 or 6.

[Claim 8] A method for producing the enzyme according to claim 1 or 2 wherein the transformant of claim 7 is cultured and the enzyme of claim 1 or 2 is collected from the culture.

[Claim 9] A protein which comprises an active domain of O-glycan α 2,8-sialyltransferase having any one of the following amino acid sequences:

- (1) an amino acid sequence corresponding to a portion between positions 26 and 398 of the amino acid sequence shown in SEQ ID NO: 1; or
- (2) an amino acid sequence comprising a deletion, substitution, and/or addition of one or several amino acids with respect to the amino acid sequence corresponding to a portion between positions 26 and 398 of the amino acid sequence shown in SEQ ID NO: 1, and having O-glycan α 2,8-sialyltransferase activity;

[Claim 10] An extracellular secretory protein, which comprises a polypeptide portion which is an active domain of the O-glycan α 2,8-sialyltransferase of claim 1 or 2, and a signal peptide, and has O-glycan α 2,8-sialyltransferase activity.

[Claim 11] A gene encoding the protein according to claim 9 or 10.

[Claim 12] A recombinant vector comprising the gene according to claim 11.

[Claim 13] The recombinant vector according to claim 12 which is an expression vector.

[Claim 14] A transformant transformed with the recombinant vector according to claim 12 or 13.

[Claim 15] A method for producing the protein according to claim 9 or 10 wherein the transformant of claim 14 is cultured and the protein of claim 9 or 10 is collected from the culture.

[Detailed Description of the Invention]

[0001]

[Industrially Applicable Field]

The present invention relates to a glycosylating enzyme and DNA encoding the enzyme. More specifically, the present invention relates to an enzyme (O-glycan α 2,8-sialyltransferase, ST8Sia VI) that efficiently transfers sialic acid through an α 2,8 linkage onto the sialic acid portion of a sugar chain having a Sia α 2,3(6)Gal (Sia: sialic acid; Gal: galactose) structure at the terminus of O-glycans such as mucin, and DNA encoding the above enzyme. The O-glycan α 2,8-sialyltransferase of the present invention is useful as a medicament having effects of suppression of cancer metastasis, prevention of virus infection, suppression of inflammatory response or activation of neural cells, as a reagent for increasing physiological action by adding sialic acid to a sugar chain, or as an enzyme inhibitor.

[0002]

[Prior Art]

Sialic acid is a substance responsible for important physiological actions such as cell-cell communication, cell-substrate interaction, and cell adhesion. The presence of sialic acid-containing sugar chains has been known, and some of such chains are expressed in stage-specific manner during development and differentiation, or in tissue-specific manner. Sialic acid exists at the terminal position of the sugar chain of a glycoprotein or glycolipid. Introduction of sialic acid into these sites is carried out enzymatically by transfer of sialic acid portion from CMP-Sia.

[0003]

Enzymes having a function in such enzymatic introduction of sialic acid (sialic acid transfer) belong to a member of glycosyltransferases called sialyltransferases. So far, 18 types of sialyltransferases have been known with regard to mammals. These sialyltransferases are broadly divided into 4 families (Tsuji, S. (1996) *J. Biochem.* 120, 1-13). This is to say, these 4 families are: α 2,3-sialyltransferase (ST3Gal-family) that transfers sialic acid onto galactose through an α 2,3 linkage; α 2,6-sialyltransferase (ST6Gal-family) that transfers sialic acid onto galactose through an α 2,6 linkage; GalNAc α 2,6-sialyltransferase (ST6GalNAc-family) that transfers sialic acid onto N-acetylgalactosamine through an α 2,6 linkage; and α 2,8-sialyltransferase (ST8Sia-family) that transfers sialic acid onto sialic acid through an α 2,8 linkage.

[0004]

Of these, with regard to α 2,8-sialyltransferase, cDNA cloning of 5 types of the enzymes (ST8Sia I-V) have been achieved so far, and their enzymatic properties have been elucidated (Yamamoto, A. *et al.* (1996) *J. Neurochem.* 66, 26-34; Kojima, N. *et al.* (1995) *FEBS Lett.* 360, 1-4; Yoshida, Y. *et al.* (1995) *J. Biol. Chem.* 270, 14628-14633; Yoshida, Y. *et al.* (1995) *J. Biochem.* 118, 658-664; Kono, M. *et al.* (1996) *J. Biol. Chem.* 271, 29366-29371). ST8Sia I is an enzyme for synthesizing a ganglioside GD3, and ST8Sia V is also an enzyme for synthesizing gangliosides GD1c, GT1a, GQ1b, GT3, and so on. ST8Sia II and IV are enzymes for synthesizing polysialic acid on the N-glycans of a neural cell adhesion molecule (NCAM). ST8Sia III is an enzyme for transferring sialic acid onto Sia α 2,3Gal β 1,4GlcNAc structures found in the N-glycans of glycoproteins and glycolipids. The preferred substrates for all of these enzymes are glycolipids or N-glycans. There have been only two reports in which these enzymes exhibit activity toward O-glycans. A case where ST8Sia II and IV synthesize oligosialic acid/polysialic acid on O-glycans found in an isoform of NCAM, and a case where ST8Sia III acts on the O-glycans of an adipocyte-specific glycoprotein AdipoQ (Suzuki, M. *et al.* (2000) *Glycobiology* 10, 1113; and Sato C, *et al.* (2001) *J. Biol.*

Chem. 276, 28849-28856). Thus, the previously reported $\alpha 2,8$ -sialyltransferases do not generally utilize O-glycans as preferred substrates. The existence of $\alpha 2,8$ -sialyltransferase which utilizes such an O-glycans as preferred substrates has been unknown.

[0005]

[Object to be Solved by the Invention]

As stated above, only 5 types of $\alpha 2,8$ -sialyltransferases have been known so far. Main substrates for all of these enzymes are glycoproteins having N-glycans or glycolipids such as gangliosides. These enzymes show no activity toward glycoproteins having O-glycans, or show only a limited activity. It is the object of the present invention to solve this problem and to provide a novel O-glycan $\alpha 2,8$ -sialyltransferase showing high activity toward O-glycans. It is also the object of the present invention to clone the cDNA encoding O-glycan $\alpha 2,8$ -sialyltransferase, so as to provide a DNA sequence encoding the above O-glycan $\alpha 2,8$ -sialyltransferase and an amino acid sequence of the above enzyme. Moreover, it is also the object of the present invention to allow a portion necessary for the activity of the above O-glycan $\alpha 2,8$ -sialyltransferase to express as a protein in a large quantity.

[0006]

[Means for Solving the Object]

The present inventors have made intensive studies to achieve the above-described objects. The present inventors have screened mouse brain and heart cDNA libraries, and have also performed PCR using cDNA derived from mouse kidney as a template, so that they have succeeded in cloning the cDNA encoding O-glycan $\alpha 2,8$ -sialyltransferase. Thus, the present invention has been completed.

That is to say, the present invention provides O-glycan $\alpha 2,8$ -sialyltransferase, which is characterized in that it has the following substrate specificity and substrate selectivity.

Substrate specificity: the substrates of the enzyme are glycoconjugates having a Sia α 2,3(6)Gal structure (wherein Sia represents sialic acid and Gal represents galactose) at the terminus thereof.

Substrate selectivity: the enzyme incorporates sialic acids into O-glycans more preferentially than into glycolipids or N-glycans.

[0007]

Preferably, the present invention provides O-glycan α 2,8-sialyltransferase having either one of the following amino acid sequences:

- (1) an amino acid sequence shown in SEQ ID NO: 1; or
- (2) an amino acid sequence comprising a deletion, substitution, and/or addition of one or several amino acids with respect to the amino acid sequence shown in SEQ ID NO: 1, and having O-glycan α 2,8-sialyltransferase activity.

[0008]

In another aspect of the present invention, the O-glycan α 2,8-sialyltransferase gene encoding the above-described amino acid sequence of the O-glycan α 2,8-sialyltransferase of the present invention is provided.

[0009]

Preferably, the present invention provides the O-glycan α 2,8-sialyltransferase gene having any one of the following nucleotide sequences:

- (1) a nucleotide sequence corresponding to a portion between nucleotide 77 and nucleotide 1270 of a nucleotide sequence shown in SEQ ID NO: 2; or
- (2) a nucleotide sequence comprising a deletion, substitution, and/or addition of one or several nucleotides with respect to the nucleotide sequence corresponding to a portion between nucleotide 77 and nucleotide 1270 of the nucleotide sequence shown in SEQ ID NO: 2, and encoding a protein having O-glycan α 2,8-sialyltransferase activity.

[0010]

In another aspect of the present invention, the followings are provided: a recombinant vector (preferably, an expression vector) comprising the above-described O-glycan α 2,8-sialyltransferase gene of the present invention; a transformant transformed with the above recombinant vector; and a method for producing the enzyme of the present invention wherein the above transformant is cultured and the enzyme of the present invention is collected from the culture.

[0011]

In another aspect of the present invention, a protein which comprises an active domain of O-glycan α 2,8-sialyltransferase having any one of the following amino acid sequences is provided:

- (1) an amino acid sequence corresponding to a portion between positions 26 and 398 of the amino acid sequence shown in SEQ ID NO: 1; or
- (2) an amino acid sequence comprising a deletion, substitution, and/or addition of one or several amino acids with respect to the amino acid sequence corresponding to a portion between positions 26 and 398 of the amino acid sequence shown in SEQ ID NO: 1, and having O-glycan α 2,8-sialyltransferase activity.

[0012]

In another aspect of the present invention, an extracellular secretory protein is provided, which comprises a polypeptide portion of the active domain and a signal peptide of the O-glycan α 2,8-sialyltransferase of the present invention, and has O-glycan α 2,8-sialyltransferase activity.

In another aspect of the present invention, a gene encoding the above-described extracellular secretory protein of the present invention is provided.

[0013]

In another aspect of the present invention, the followings are provided: a recombinant vector (preferably, an expression vector) comprising a gene encoding the above-described extracellular secretory protein of the present invention; a transformant

transformed with the above recombinant vector; and a method for producing the protein of the present invention wherein the above transformant is cultured and the enzyme of the present invention is collected from the culture.

[0014]

[Mode for Carrying out the Invention]

The embodiments of the present invention and the methods for carrying out the present invention will be described in detail below.

(1) Enzyme and protein of the present invention

The O-glycan α 2,8-sialyltransferase of the present invention is characterized in that it has the following substrate specificity and substrate selectivity.

Substrate specificity: the substrates of the enzyme are glycoconjugates having a Sia α 2,3(6)Gal structure (wherein Sia represents sialic acid and Gal represents galactose) at the terminus thereof.

Substrate selectivity: the enzyme incorporates sialic acids into O-glycan more preferentially than into glycolipids or N-glycans.

[0015]

The above-described substrate specificity and substrate selectivity are characteristics which have been demonstrated by mouse-derived O-glycan α 2,8-sialyltransferases obtained in examples described in the present specification. The O-glycan α 2,8-sialyltransferase of the present invention is not only derived from a mouse, and it is easily understandable for a person skilled in the art that the same type of O-glycan α 2,8-sialyltransferase exists in the tissues of other mammals and that those O-glycan α 2,8-sialyltransferases have a high homology to one another.

[0016]

Such O-glycan α 2,8-sialyltransferases are characterized in that they have the above-described substrate specificity and substrate selectivity. These enzymes are also included in the scope of the present invention.

Examples of such an O-glycan α 2,8-sialyltransferase may include natural enzymes derived from mammalian tissues and mutants thereof, and extracellular secretory proteins catalyzing the transfer of sialic acid to O-glycans through an α 2,8-linkage, which are produced by genetic recombination, such as those produced in examples described later. These are also included in the scope of the present invention.

[0017]

O-glycan α 2,8-sialyltransferase having either one of the following amino acid sequences may be one example of the O-glycan α 2,8-sialyltransferase of the present invention:

- (1) an amino acid sequence shown in SEQ ID NO: 1; or
- (2) an amino acid sequence comprising a deletion, substitution, and/or addition of one or several amino acids with respect to the amino acid sequence shown in SEQ ID NO: 1, and having O-glycan α 2,8-sialyltransferase activity.

[0018]

In addition, it is to be understood that an active domain of the O-glycan α 2,8-sialyltransferase of the present invention and proteins having O-glycan α 2,8-sialyltransferase activity obtained by alteration or modification of a portion of the amino acid sequence thereof are all included in the scope of the present invention. Preferred examples of such an active domain may include an active domain of O-glycan α 2,8-sialyltransferase corresponding to a portion between positions 26 and 398 of the amino acid sequence shown in SEQ ID NO: 1. A sequence portion between positions 26 and approximately 100 of the amino acid sequence shown in SEQ ID NO: 1 is a region called stem, and it is considered that this region is not necessarily required for the activity. Accordingly, a region corresponding to positions 101 to 398 of the amino acid sequence shown in SEQ ID NO: 1 may be used as an active domain of O-glycan α 2,8-sialyltransferase.

[0019]

That is to say, the present invention provides a protein which comprises an active domain of O-glycan α 2,8-sialyltransferase having any one of the following amino acid sequences:

- (1) an amino acid sequence corresponding to a portion between positions 26 and 398 of the amino acid sequence shown in SEQ ID NO: 1; or
- (2) an amino acid sequence comprising a deletion, substitution, and/or addition of one or several amino acids with respect to the amino acid sequence corresponding to a portion between positions 26 and 398 of the amino acid sequence shown in SEQ ID NO: 1, and having O-glycan α 2,8-sialyltransferase activity.

[0020]

In the present specification, the range of "one or several" in the expression "an amino acid sequence comprising a deletion, substitution, and/or addition of one or several amino acids" is not particularly limited. For example, it means 1 to 20 amino acids, preferably 1 to 10 amino acids, more preferably 1 to 7 amino acids, further more preferably 1 to 5 amino acids, and particularly preferably 1 to 3 amino acids.

[0021]

A method for obtaining the enzyme or protein of the present invention is not particularly limited. The protein of the present invention may be a protein synthesized by chemical synthesis, or recombinant protein produced by genetic recombination.

When a recombinant protein is produced, first, DNA encoding the protein is required to be obtained. Suitable primers are designed based on the information regarding amino acid sequence shown in SEQ ID NO: 1 and nucleotide sequence shown in SEQ ID NO: 2 of the sequence listing in the present specification. Thereafter, using the obtained primers, PCR is carried out with a suitable cDNA library as a template, so as to obtain DNA encoding the enzyme of the present invention.

For example, a method for isolating cDNA encoding O-glycan α 2,8-sialyltransferases having amino acid sequences shown in SEQ ID NO: 1 is

described in detail in examples described later. However, a method for isolating cDNA encoding the O-glycan α 2,8-sialyltransferase of the present invention is not limited thereto. A person skilled in the art could easily isolate cDNA of interest by referring to the methods described in examples below and appropriately modifying or altering them.

[0022]

Moreover, when a partial fragment of DNA encoding the enzyme of the present invention is produced by the above-described PCR, the produced DNA fragments can be successively ligated to one another, so as to obtain DNA encoding a desired enzyme. The obtained DNA can be then introduced into a suitable expression system, so as to generate the enzyme of the present invention. Expression of the enzyme in such an expression system will be described later in the specification.

[0023]

An extracellular secretory protein, which comprises a polypeptide portion of the active domain of the O-glycan α 2,8-sialyltransferase of the present invention and a signal peptide, and has O-glycan α 2,8-sialyltransferase activity is also included in the present invention.

[0024]

In some cases, the O-glycan α 2,8-sialyltransferase of the present invention may remain in cells after the expression and may not be secreted outside of the cells. In addition, there is a possibility that the production of the enzymes may be decreased when the intracellular concentration thereof exceeds a certain limit. In order to effectively use the activity of the above O-glycan α 2,8-sialyltransferase to transfer sialic acid to O-glycans through an α 2,8-linkage, a soluble form of proteins retaining the activities of the present enzymes and being secreted from cells during the expression may be produced. An example of such a protein may be an extracellular secretory protein, which comprises a signal peptide and a polypeptide portion of the active domain of O-glycan α 2,8-sialyltransferase which is involved in the activity of the O-glycan

α 2,8-sialyltransferase of the present invention, and catalyzes the transfer of sialic acid to O-glycans through an α 2,8-linkage. For example, a fusion protein with protein A as described in the specification is preferred embodiments of the secretory protein of the present invention.

[0025]

Sialyltransferases that have been cloned so far have a domain structure similar to that of other glycosyltransferases. This is to say, the previously cloned sialyltransferases comprise an NH_2 -terminal short cytoplasmic tail, a hydrophobic signal anchor domain, a stem region having proteolytic sensitivity, and a COOH-terminal large active domain (Paulson, J.C. and Colley, K.J., *J. Biol. Chem.*, 264, 17615-17618, 1989). In order to examine the position of a transmembrane domain of the O-glycan α 2,8-sialyltransferase of the present invention, a hydropathy plot prepared according to the method of Kyte and Doolittle (Kyte, J. and Doolittle, R.F., *J. Mol. Biol.*, 157, 105-132, 1982) can be used. Moreover, in order to estimate an active domain portion, recombinant plasmids into which various types of fragments are introduced are produced and used. An example of such methods is described in detail, for example, in PCT/JP94/02182. However, a method for confirming the position of a transmembrane domain or estimating an active domain portion is not limited thereto.

[0026]

In order to produce an extracellular secretory protein which comprises a polypeptide portion of the active domain of O-glycan α 2,8-sialyltransferase and a signal peptide, for example, a sequence corresponding to the active domain of O-glycan α 2,8-sialyltransferase may be subjected to inframe fusion with an immunoglobulin signal peptide sequence as a signal peptide. As such a method, the method of Jobling (Jobling, S.A. and Gehrke, L., *Nature* (Lond.), 325, 622-625, 1987), for example, can be used. Further, as is described in detail in examples of the present specification, a fusion protein with protein A may also be produced. However, the type of a signal peptide, the

method of the fusion of a signal peptide with an active domain, and the method of solubilization are not limited to those described above. A person skilled in the art may appropriately select a polypeptide portion which is an active domain of O-glycan α 2,8-sialyltransferase, and may fuse the selected polypeptide portion with any available signal peptide by a suitable method, so as to produce an extracellular secretory protein.

[0027]

(2) Gene of the present invention

The present invention provides a gene encoding the amino acid sequence of the O-glycan α 2,8-sialyltransferase of the present invention.

Specific examples of a gene encoding the amino acid sequence of the O-glycan α 2,8-sialyltransferase of the present invention may include genes having any one of the following nucleotide sequences:

- (1) a nucleotide sequence corresponding to a portion between nucleotide 77 and nucleotide 1270 of a nucleotide sequence shown in SEQ ID NO: 2; or
- (2) a nucleotide sequence comprising a deletion, substitution, and/or addition of one or several nucleotides with respect to the nucleotide sequence corresponding to a portion between nucleotide 77 and nucleotide 1270 of the nucleotide sequence shown in SEQ ID NO: 2, and encoding a protein having O-glycan α 2,8-sialyltransferase activity.

[0028]

The range of “one or several” in the expression “a nucleotide sequence comprising a deletion, substitution, and/or addition of one or several nucleotides” in the present specification is not particularly limited. For example, it means 1 to 60 nucleotides, preferably 1 to 30 nucleotides, more preferably 1 to 20 nucleotides, further more preferably 1 to 10 nucleotides, further more preferably 1 to 5 nucleotides, and particularly preferably 1 to 3 nucleotides.

[0029]

A gene encoding a protein comprising an active domain of the O-glycan α 2,8-sialyltransferase of the present invention, and a gene encoding an extracellular secretory protein which comprises a polypeptide portion which is the above active domain and a signal peptide and has O-glycan α 2,8-sialyltransferase activity, are also included in the scope of the present invention.

[0030]

The gene of the present invention can be obtained by the above-described method.

A method of introducing a desired mutation into a certain nucleic acid sequence is known to those skilled in the art. For example, known techniques such as site-directed mutagenesis, PCR using degenerated oligonucleotides, or exposure of cells containing nucleic acid to a mutagenic agent or radioactive ray are used as appropriate, whereby DNA comprising a mutation can be constructed. Such known techniques are described, for example, in Molecular Cloning: A laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY., 1989; and Current Protocols in Molecular Biology, Supplements 1 to 38, John Wiley & Sons (1987-1997).

[0031]

(3) Recombinant vector of the present invention

The gene of the present invention can be inserted into a suitable vector and used. The type of a vector used in the present invention is not particularly limited. For example, it may be autonomously replicating vector (e.g., a plasmid, etc.), or it may be a vector which is incorporated into the genome in host cells when it is introduced into the host cells, and replicates with an incorporated chromosome.

The vector used in the present invention is preferably an expression vector. In an expression vector, elements necessary for transcription (e.g., a promoter, etc.) are functionally ligated to the gene of the present invention. A promoter is a DNA sequence having transcription activity in host cells, and it can appropriately be selected depending on the type of host cells.

[0032]

Examples of a promoter capable of functioning in bacterial cells may include a *Bacillus stearothermophilus* maltogenic amylase gene promoter, a *Bacillus licheniformis* alpha-amylase gene promoter, a *Bacillus amyloliquefaciens* BAN amylase gene promoter, a *Bacillus subtilis* alkaline protease gene promoter, a *Bacillus pumilus* xylosidase gene promoter, a phage λ P_R or P_L promoter, and an *Escherichia coli* lac, trp, or lac promoter.

[0033]

Examples of a promoter capable of functioning in mammalian cells may include an SV40 promoter, an MT-1 (metallothionein gene) promoter, and an adenovirus 2 major late promoter. Examples of a promoter capable of functioning in insect cells may include a polyhedrin promoter, a P10 promoter, an *Autographa californica* polyhedrosis basic protein promoter, a baculovirus immediate early gene 1 promoter, and a baculovirus 39K delayed-early gene promoter. Examples of a promoter capable of functioning in yeast host cells may include a promoter derived from a yeast glycolytic system gene, an alcohol dehydrogenase gene promoter, a TPI1 promoter, and an ADH2-4c promoter.

Examples of a promoter capable of functioning in filamentous cells may include an ADH3 promoter and a tpiA promoter.

[0034]

The DNA of the present invention may be functionally ligated to a human growth hormone terminator, or in the case where a host is Mycomycete, the DNA may be functionally ligated to an appropriate terminator such as a TPI1 terminator or ADH3 terminator, as necessary. The recombinant vector of the present invention may also comprise elements such as a polyadenylation signal (e.g., those derived from SV40 or adenovirus 5E1b region), a transcription enhancer sequence (e.g., SV40 enhancer), and a translation enhancer sequence (e.g., those encoding adenovirus VA RNA).

The recombinant vector of the present invention may further comprise a DNA sequence enabling the vector to replicate in host cells. An example may include an SV40 replication origin (when the host cells are mammalian cells).

[0035]

The recombinant vector of the present invention may further comprise a selective marker. Examples of a selective marker may include genes whose complements are deficient in host cells, such as dihydrofolate reductase (DHFR) or a *Schizosaccharomyces pombe* TPI gene, and drug resistant genes that are resistant to ampicillin, kanamycin, tetracycline, chloramphenicol, neomycin, hygromycin, etc.

A method of ligating the DNA of the present invention, a promoter, and a terminator and/or a secretory signal sequence, as desired, to one another, and inserting them into a suitable vector has been well known to those skilled in the art.

[0036]

(4) Transformant of the present invention, and production of protein using the same

The DNA or recombinant vector of the present invention can be introduced into a suitable host, so as to prepare a transformant.

Any cells may be used as host cells into which the DNA or recombinant vector of the present invention is introduced, as long as they allow the DNA construct of the present invention to express therein. Examples of host cells may include bacteria, yeasts, Mycomycetes, and higher eukaryotes.

[0037]

Examples of bacterial cells may include Gram-positive bacteria such as *Bacillus* or *Streptomyces*, and Gram-negative bacteria such as *Escherichia coli*. Transformation of these bacteria may be carried out by the protoplast method or known methods, using competent cells.

Examples of mammalian cells may include HEK293 cells, HeLa cells, COS cells, BHK cells, CHL cells, and CHO cells. A method of transforming mammalian cells and

allowing a DNA sequence introduced into the cells to express therein has also been known. Examples of such a method may include the electroporation, the calcium phosphate method, and the lipofection method.

[0038]

Examples of yeast cells may include cells belonging to *Saccharomyces* or *Schizosaccharomyces*. Examples of such cells may include *Saccharomyces cerevisiae* and *Saccharomyces kluyveri*. Examples of a method of introducing a recombinant vector into a yeast host may include the electroporation, the spheroplast method, and the lithium acetate method.

[0039]

Examples of other fungal cells may include cells belonging to filamentous fungi such as *Aspergillus*, *Neurospora*, *Fusarium*, or *Trichoderma*. When filamentous fungi are used as host cells, transformation can be carried out by incorporating a DNA construct into a host chromosome to obtain recombinant host cells. Such a DNA construct can be incorporated into a host chromosome according to known methods such as homologous recombination or heterologous recombination.

[0040]

When insect cells are used as host cells, a recombinant gene-introduced vector and baculovirus are co-introduced into insect cells, and recombinant virus is obtained in the culture supernatant of the insect cells. Thereafter, insect cells are infected with the recombinant virus, so that a protein is expressed (which is described in e.g. *Baculovirus Expression Vectors, A Laboratory Manual*; and *Current Protocols in Molecular Biology, Bio/Technology*, 6, 47 (1998)).

[0041]

As an example of baculovirus, *Autographa californica* nuclear polyhedrosis virus infecting *Manestra* insects can be used.

Examples of insect cells used herein may include *Spodoptera frugiperda* ovarian cells Sf 9 and Sf21 [Baculovirus Expression Vectors, A Laboratory Manual, W. H. Freeman and Company, New York (1992)], and *Trichoplusia ni* ovarian cells HiFive (manufactured by Invitrogen).

Examples of a method of co-introducing a recombinant gene-introduced vector and the above baculovirus into insect cells to prepare recombinant virus may include the calcium phosphate method and the lipofection method.

[0042]

The above transformant is cultured in a nutrient medium under conditions enabling the expression of the introduced DNA construct. In order to isolate and purify the enzyme of the present invention from the culture of the transformant, common protein isolation and purification methods may be applied.

For example, where the enzyme of the present invention is expressed in a state where it is dissolved in cells, the cells are recovered by centrifugation after completion of the culture, and they are then suspended in a water-type buffer solution. Thereafter, the cells were disintegrated with an ultrasonic disintegrator or the like, so as to obtain a cell-free extract. A purified sample can be obtained from a supernatant obtained by centrifuging the above cell-free extract, using singly or in combination the following common protein isolation and purification methods: solvent extraction method, salting-out using ammonium sulfate or the like, desalting, precipitation method using organic solvents, anion exchange chromatography using resin such as diethylaminoethyl (DEAE) sepharose, cation exchange chromatography using resin such as S-Sepharose FF (manufactured by Pharmacia), hydrophobic chromatography using resin such as butyl sepharose or phenyl sepharose, gel filtration using a molecular sieve, affinity chromatography, chromatofocusing, electrophoresis such as isoelectric focusing, etc.

The present invention will be further specifically described in the following examples. However, these examples are not intended to limit the scope of the present invention.

[0043]

[Examples]

The following reagents and samples were used in specific examples of the present invention. Fetuin, asialofetuin, bovine submaxillary mucin (BSM), α 1-acid glycoprotein, ovomucoid, lactosyl ceramide (LacCer), GM3, GM1a, GD1a, GD1b, GT1b, CMP-NeuAc, 6'-sialyllactose, 3'-sialyl-*N*-acetylglucosamine, and Triton CF-54 were purchased from Sigma. 3'-sialyllactose and 6'-sialyl-*N*-acetylglucosamine were purchased from Calbiochem. *N*-acetylneuraminic acid (NeuAc), GM4, Gal, and *N*-acetylglucosamine (GalNAc) were purchased from Wako Pure Chemical Industries, Ltd. GD3 was purchased from Snow Brand Milk Products Co., Ltd. GQ1b was purchased from Alexis Biochemicals. CMP-[14 C]-NeuAc (12.0 GBq/mmol) was purchased from Amersham Pharmacia Biotech. Sialidases (NANase II, III) were purchased from Glyko Inc. *N*-glycanase (Glycopeptidase F) was purchased from Takara Shuzo Co., Ltd. [α - 32 P]dCTP was purchased from NEN. GM1b and its positional analogs, GSC-68, 2,3-sialylparagloboside (2,3-SPG), and 2,6-sialylparagloboside (2,6-SPG) were contributed from Prof. Makoto Kiso (Faculty of Agriculture, Gifu University). NeuAc α 2,3Gal and NeuAc α 2,6Gal were contributed from Dr. Hideki Ishida (The Noguchi Institute). An anti-GD3 monoclonal antibody KM641 was contributed from Dr. Kenya Shitara and Dr. Nobuo Hanai of Kyowa Hakko Kogyo Co., Ltd. Peroxidase-conjugated AffiniPure goat anti-mouse IgG + IgM (H + L) was purchased from Jackson Immuno Research. Desialylated (asialo) glycoproteins obtained by removing sialic acids from BSM, α 1-acid glycoprotein, and ovomucoid were prepared by treating them at 80°C for 1 hour in 0.02 N HCl.

[0044]

Using the amino acid sequence of mouse sialyltransferase ST8Sia V, a clone encoding a novel sialyltransferase showing a homology with the above enzyme has been searched against the database of expressed sequence tag (dbEST) of the National Center for Biotechnology Information. As a result, clones deposited under GenBankTM accession Nos. BE633149, BE686184, and BF730564 were obtained. Based on the information regarding the nucleotide sequences of these clones, two types of synthetic DNA fragments, 5'-CTTTCTGGAGAACTAAAGG-3' (corresponding to nucleotides 1001-1020 in Figure 1) (SEQ ID NO: 3) and 5'-AATTGCAGTTTGAGGATTCC-3' (corresponding to a complementary strand of nucleotides 1232-1251 in Figure 1) (SEQ ID NO: 4) were prepared. Thereafter, in accordance with the method of Israel (Israel, D. I. (1993) *Nucleic Acids Res.* 21, 2627-2631), the cDNA library of each of mouse brain and heart was screened by the polymerase chain reaction method (PCR). As a result, a clone encoding a portion of a novel sialyltransferase was obtained from each cDNA library. In order to obtain a full-length clone, two types of synthetic DNA fragments 5'-TGGCTCAGGATGAGATCGGG-3' (corresponding to nucleotides 68-87 in Figure 1) (SEQ ID NO: 5) and 5'-TACTAGCGCTCCCTGTGATTGG-3' (corresponding to a complementary strand of nucleotides 725-746 in Figure 1) (SEQ ID NO: 6) were further prepared. Thereafter, using mouse kidney-derived cDNA as a template, DNA located between both the synthetic DNA fragments was amplified by PCR. The amplified fragment was ligated to a clone obtained from the mouse brain cDNA library, so as to obtain a full-length clone. This cDNA had a single open reading frame encoding type II transmembrane protein of 398 amino acids with an estimated molecular weight of 45,399. In addition, sialyl motifs conserved in sialyltransferases were present in the amino acid sequence thereof. This protein showed 42.0% and 38.3% homology with ST8Sia I and V, respectively, at an amino acid sequence level among known mouse sialyltransferases (Figure 2). As described below, since this protein had the activity of

α 2,8-sialyltransferase, it was named as the *O*-glycan α 2,8-sialyltransferase of the present invention, ST8Sia VI.

[0045]

In order to examine enzymatic properties of the protein, a secretory protein was produced. First, with regard to mouse ST8Sia VI, using two types of synthetic DNA fragments each containing a *Xho*I site, 5'-TGCTCTCGAGCCCAGCCGACGCGCCTGCCC-3' (corresponding to nucleotides 141-170 in Figure 1) (SEQ ID NO: 7) and 5'-TATTCTCGAGCTAAGAAACGTTAAGCCGTT-3' (corresponding to a complementary strand of nucleotides 1263-1293 in Figure 1) (SEQ ID NO: 8), a DNA fragment encoding the active domain of mouse ST8Sia VI was amplified by PCR with cloned full-length cDNA as a template. The amplified product was cleaved with *Xho*I, and a cleaved portion was inserted into the *Xho*I site of a mammalian expression vector, pcDSA. The obtained expression vector was named as pcDSA-ST8Sia VI. This encodes a secretory fusion protein comprising a signal peptide of mouse immunoglobulin IgM, *Staphylococcus aureus* protein A, and the active domain of ST8Sia VI. Using pcDSA-ST8Sia VI and lipofectamine (Invitrogen), transient expression was carried out in COS-7 cells (Kojima, N. *et al.* (1995) *FEBS Lett.* **360**, 1-4). The protein of the present invention secreted from the cell into which pcDSA-ST8Sia VI had been introduced was named as PA-ST8Sia VI. PA-ST8Sia VI was adsorbed to IgG-Sepharose (Amersham Pharmacia Biotech), and were then recovered from medium.

[0046]

Sialyltransferase activity was measured as follows according to the method of Lee *et al.* (Lee, Y.-C. *et al.* (1999) *J. Biol. Chem.* **274**, 11958-11967). A reaction solution (10 μ l) containing 50 mM MES buffer (pH 6.0), 1 mM MgCl₂, 1 mM CaCl₂, 0.5% Triton CF-54, 100 μ M CMP-[¹⁴C]-NeuAc, a glycoconjugate (which was added at 0.5 mg/ml in the case of glycolipids, and at 1 mg/ml in the case of glycoproteins or oligosaccharides),

and a PA-ST8Sia VI suspension, was incubated at 37°C for 3 to 20 hours. Thereafter, in the case of glycolipids, the reaction product was purified with a C-18 column (Sep-Pak Vac 100 mg; Waters) and the purified product was used as a sample, and in the case of oligosaccharides or glycoproteins, the reaction product was directly used as a sample. Thus, the obtained samples were subjected to analysis. In the case of oligosaccharides or glycolipids, the sample was spotted on a silica gel 60 HPTLC plate (Merck), and was then developed with a developing solvent consisting of ethanol : pyridine : n-butanol : water : acetic acid = 100 : 10 : 10 : 30 : 3 (for oligosaccharides), or a developing solvent consisting of chloroform : methanol : 0.02% CaCl₂ = 55 : 45 : 10 (for glycolipids). In the case of glycoproteins, analysis was carried out by SDS-polyacrylamide gel electrophoresis. The obtained radioactivities were visualized with a BAS2000 radio image analyzer (Fuji Film) and then quantified.

Table 1 shows substrate specificity of PA-ST8Sia VI.

[0047]

[Table 1]

Table 1. Acceptor substrate specificity of ST3Sia VI. Using PNA-ST3Sia VI, specificity against various acceptor substrates was examined. The concentration of U-6 substrates is 0.5 mg/ml in the case of glycosylated, and 1 mg/ml in the case of glycoproteins, monosaccharides and oligosaccharides. The relative activity was calculated by taking incorporation obtained with FcRan (2.08 pmol/h/ml enzyme solution) as 100. R represents the remainder of the N-linked sugar chain.

Acceptor substrate	Representative structure	Relative activity (%)
Glycoproteins		
FcRan	NeuAc2,3Galβ1,3GalNAc-O-SerThr NeuAc2,3Galβ1,3NeuAc2,6GalNAc-O-SerThr NeuAc2,6Galβ1,4GlcNAc-R	100
AsialofcRan	NeuAc2,6Galβ1,4GlcNAc-R	0
α1-Acid glycoprotein	NeuAc2,6Galβ1,4GlcNAc-R	0
Asialbo-α1-Acid glycoprotein	NeuAc2,6Galβ1,4GlcNAc-R	0
BSM	NeuAc2,6GalNAc-O-SerThr GlcNAcβ1,3NeuAc2,6GalNAc-O-SerThr	375
Asialbo-BSM	NeuAc2,6GalNAc-O-SerThr	0
Ovomucoid	NeuAc2,3Galβ1,4GlcNAc-R	62
Asialbo-ovomucoid	NeuAc2,3Galβ1,4GlcNAc-R	0
glycolipids		
Lactosylceramide	Galβ1,4Glcβ1-Cer	0
GM4	NeuAc2,3Galβ1-Cer	1.0
GM3	NeuAc2,3Galβ1,4Glcβ1-Cer	13.0
GM1a	Galβ1,3GalNAcβ1,4NeuAc2,3Galβ1,4Glcβ1-Cer	0
GM1b	NeuAc2,3Galβ1,3GalNAcβ1,4NeuAc2,3Galβ1,4Glcβ1-Cer	6.0
GM2	NeuAc2,8NeuAc2,3Galβ1,4Glcβ1-Cer	0
GM3	NeuAc2,8NeuAc2,3Galβ1,4Glcβ1-Cer	0
GM1b	NeuAc2,8NeuAc2,3Galβ1,4Glcβ1-Cer	1.1
GM1c	NeuAc2,8NeuAc2,3Galβ1,4Glcβ1-Cer	0
GM1d	NeuAc2,8NeuAc2,3Galβ1,4Glcβ1-Cer	1.0
GM1e	NeuAc2,8NeuAc2,3Galβ1,4Glcβ1-Cer	2.6
GM1f	NeuAc2,8NeuAc2,3Galβ1,4Glcβ1-Cer	3.6
GM1g	NeuAc2,8NeuAc2,3Galβ1,4Glcβ1-Cer	3.6
GM1h	NeuAc2,8NeuAc2,3Galβ1,4Glcβ1-Cer	3.6
GM1i	NeuAc2,8NeuAc2,3Galβ1,4Glcβ1-Cer	3.6
GM1j	NeuAc2,8NeuAc2,3Galβ1,4Glcβ1-Cer	3.6
GM1k	NeuAc2,8NeuAc2,3Galβ1,4Glcβ1-Cer	3.6
GM1l	NeuAc2,8NeuAc2,3Galβ1,4Glcβ1-Cer	3.6
GM1m	NeuAc2,8NeuAc2,3Galβ1,4Glcβ1-Cer	3.6
GM1n	NeuAc2,8NeuAc2,3Galβ1,4Glcβ1-Cer	3.6
GM1o	NeuAc2,8NeuAc2,3Galβ1,4Glcβ1-Cer	3.6
GM1p	NeuAc2,8NeuAc2,3Galβ1,4Glcβ1-Cer	3.6
GM1q	NeuAc2,8NeuAc2,3Galβ1,4Glcβ1-Cer	3.6
GM1r	NeuAc2,8NeuAc2,3Galβ1,4Glcβ1-Cer	3.6
GM1s	NeuAc2,8NeuAc2,3Galβ1,4Glcβ1-Cer	3.6
GM1t	NeuAc2,8NeuAc2,3Galβ1,4Glcβ1-Cer	3.6
GM1u	NeuAc2,8NeuAc2,3Galβ1,4Glcβ1-Cer	3.6
GM1v	NeuAc2,8NeuAc2,3Galβ1,4Glcβ1-Cer	3.6
GM1w	NeuAc2,8NeuAc2,3Galβ1,4Glcβ1-Cer	3.6
GM1x	NeuAc2,8NeuAc2,3Galβ1,4Glcβ1-Cer	3.6
GM1y	NeuAc2,8NeuAc2,3Galβ1,4Glcβ1-Cer	3.6
GM1z	NeuAc2,8NeuAc2,3Galβ1,4Glcβ1-Cer	3.6
monosaccharides and oligosaccharides		
3'-Sialyllactose	NeuAc2,3Galβ1,4Glc	629
6'-Sialyllactose	NeuAc2,3Galβ1,4Glc	91.5
3'-Sialyl-N-acetylglucosamine	NeuAc2,3Galβ1,4GlcNAc	411
6'-Sialyl-N-acetylglucosamine	NeuAc2,3Galβ1,4GlcNAc	88.7
3'-Sialylgalactose	NeuAc2,3Galβ1,4Glc	13.9
6'-Sialylgalactose	NeuAc2,3Galβ1,4Glc	2.0
N-Acetylglucosamine acid	GlcNAc	0
Galactose	Glc	0
N-Acetylglucosamine	GlcNAc	0

[0048]

PA-ST8Sia VI showed activity on glycolipids having a structure "NeuAc α 2,3(6)Gal-" at the nonreducing end thereof, such as GM4, GM3, GD1a, GT1b, GM1b, GSC-68, 2,3-SPG, or 2,6-SPG. When GM3 was used as a substrate, the incorporated sialic acid of the reaction product was not cleaved with sialidase (NANase II), which specifically cleaves α 2,3- and α 2,6-linked sialic acid. However, the incorporated sialic acid was cleaved with sialidase (NANase III), which specifically cleaves α 2,3-, α 2,6-, α 2,8- and α 2,9-linked sialic acids (Figure 3A). In addition, it was confirmed by TLC immunostaining using an anti-GD3 monoclonal antibody KM641 (Saito, M. *et al.* (2000) *Biochim. Biophys. Acta* **1523**, 230-235) that this reaction product was GD3 into which sialic acid had been introduced through an α 2,8-linkage (Figure 3B). Thus, it was clarified that PA-ST8Sia VI transfers sialic acid through an α 2,8-linkage.

[0049]

On the other hand, where a glycoprotein was used as a substrate (Table 1), PA-ST8Sia showed the highest activity toward BSM, which contains only *O*-glycans as glycoconjugate. PA-ST8Sia also showed activity toward Fetuin, which contains both *O*-glycans and *N*-glycans and toward Ovomucoid, which contains only *N*-glycans. However, the activity toward Ovomucoid was lower than that toward a protein containing *O*-glycans. Moreover, PA-ST8Sia VI showed no activity on asialoglycoproteins. Furthermore, from an experiment wherein monosaccharide or oligosaccharide was used as a substrate (Table 1), it was found that the minimum sugar chain unit, which was recognized by PA-ST8Sia VI as a substrate, is NeuAc α 2,3(6)Gal.

[0050]

It was found by an *N*-glycanase treatment that when Fetuin was used as a substrate, the majority of sialic acid, which was newly introduced by PA-ST8Sia VI, was incorporated into *O*-glycans (Figure 4). That is, when Fetuin was sialylated by PA-ST8Sia VI with [14 C]-NeuAc, and the sialylated product was then treated with

N-glycanase, which releases *N*-glycans from a peptide portion. The majority (82.7%) of radioactivity was still kept in the Fetuin after this treatment. This fact shows that the majority of sialic acid introduced by PA-ST8Sia VI was incorporated into *O*-glycans. On the other hand, the same experiment was carried out using ST8Sia III which used *N*-glycans as substrates. As a result, it was found that radioactivity completely disappeared.

[0051]

Moreover, in order to clarify the substrate specificity and substrate selectivity of PA-ST8Sia VI, the K_m and V_{max} values for BSM and GM3, respectively, were obtained. With regard to BSM, the K_m value was 0.03 mM, the V_{max} value was 23.8 pmol/h/ml enzyme solution, and the V_{max}/K_m value was 793. With regard to GM3, the K_m value was 0.5 mM, the V_{max} value was 0.67 pmol/h/ml enzyme solution, and the V_{max}/K_m value was 1.34. These results show that, for PA-ST8Sia VI, *O*-glycans are much more preferable substrates than glycolipids or *N*-glycans. Accordingly, it can be said that ST8Sia VI had substrate specificity different from that of the conventional $\alpha 2,8$ -sialyltransferases.

[0052]

[Effect of the Invention]

The present invention provides a novel enzyme *O*-glycan $\alpha 2,8$ -sialyltransferase, and a novel protein having an active portion of the enzyme and being extracellularly secreted. The enzyme and protein of the present invention have the activity of *O*-glycan $\alpha 2,8$ -sialyltransferase. Accordingly, it is useful as a reagent for introducing a human-type sugar chain into a protein, for example. In addition, the *O*-glycan $\alpha 2,8$ -sialyltransferase of the present invention is useful also as a medicament for treating hereditary diseases caused by deficiency of sugar chains specific for humans. Moreover, the *O*-glycan $\alpha 2,8$ -sialyltransferase of the present invention can also be used as a medicament which acts for suppression of cancer metastasis, prevention of virus

infection, suppression of inflammatory response, or activation of neural cells. Furthermore, the O-glycan α 2,8-sialyltransferase of the present invention is useful also as a reagent used in studies for increasing physiological action by adding sialic acid to drugs or the like.

[0053]

[SEQUENCE LISTING]

SEQUENCE LISTING

<110> RIKEN

<120> Sugar chain synthetase

<130> A21046A

<160> 8

[0054]

<210> 1

<211> 398

<212> PRT

<213> Mouse

<400> 1

Met Arg Ser Gly Gly Thr Leu Phe Ala Leu Ile Gly Ser Leu Met Leu

1 5 10 15

Leu Leu Leu Leu Arg Met Leu Trp Cys Pro Ala Asp Ala Pro Ala Arg

20 25 30

Ser Arg Leu Leu Met Glu Gly Ser Arg Glu Asp Thr Ser Gly Thr Ser

35 40 45

Ala Ala Leu Lys Thr Leu Trp Ser Pro Thr Thr Pro Val Pro Arg Thr

50 55 60

Arg Asn Ser Thr Tyr Leu Asp Glu Lys Thr Thr Gln Ile Thr Glu Lys

65 70 75 80

Cys Lys Asp Leu Gln Tyr Ser Leu Asn Ser Leu Ser Asn Lys Thr Arg

85 90 95

Arg Tyr Ser Glu Asp Asp Tyr Leu Gln Thr Ile Thr Asn Ile Gln Arg

100 105 110

Cys Pro Trp Asn Arg Gln Ala Glu Glu Tyr Asp Asn Phe Arg Ala Lys
 115 120 125
 Leu Ala Ser Cys Cys Asp Ala Ile Gln Asp Phe Val Val Ser Gln Asn
 130 135 140
 Asn Thr Pro Val Gly Thr Asn Met Ser Tyr Glu Val Glu Ser Lys Lys
 145 150 155 160
 His Ile Pro Ile Arg Glu Asn Ile Phe His Met Phe Pro Val Ser Gln
 165 170 175
 Pro Phe Val Asp Tyr Pro Tyr Asn Gln Cys Ala Val Val Gly Asn Gly
 180 185 190
 Gly Ile Leu Asn Lys Ser Leu Cys Gly Ala Glu Ile Asp Lys Ser Asp
 195 200 205
 Phe Val Phe Arg Cys Asn Leu Pro Pro Ile Thr Gly Ser Ala Ser Lys
 210 215 220
 Asp Val Gly Ser Lys Thr Asn Leu Val Thr Val Asn Pro Ser Ile Ile
 225 230 235 240
 Thr Leu Lys Tyr Gln Asn Leu Lys Glu Lys Lys Ala Gln Phe Leu Glu
 245 250 255
 Asp Ile Ser Thr Tyr Gly Asp Ala Phe Leu Leu Leu Pro Ala Phe Ser
 260 265 270
 Tyr Arg Ala Asn Thr Gly Ile Ser Phe Lys Val Tyr Gln Thr Leu Lys
 275 280 285
 Glu Ser Lys Met Arg Gln Lys Val Leu Phe Phe His Pro Arg Tyr Leu
 290 295 300
 Arg His Leu Ala Leu Phe Trp Arg Thr Lys Gly Val Thr Ala Tyr Arg
 305 310 315 320
 Leu Ser Thr Gly Leu Met Ile Ala Ser Val Ala Val Glu Leu Cys Glu

325	330	335
Asn Val Lys Leu Tyr Gly Phe Trp Pro Phe Ser Lys Thr Ile Glu Asp		
340	345	350
Thr Pro Leu Ser His His Tyr Tyr Asp Asn Met Leu Pro Lys His Gly		
355	360	365
Phe His Gln Met Pro Lys Glu Tyr Ser Gln Met Leu Gln Leu His Met		
370	375	380
Arg Gly Ile Leu Lys Leu Gln Phe Ser Lys Cys Glu Thr Ala		
385	390	395
[0055]		
<210> 2		
<211> 3166		
<212> DNA		
<213> Mouse		
<400> 2		
cggagcggcg agtcgggtgcc gcccgggctg cgttcgcccc cggcagcttt ggccggcgagg	60	
acgcccgctgg ctccagg atg aga tcg ggg ggc acg ctg ttc gcc ctc ata	109	
Met Arg Ser Gly Gly Thr Leu Phe Ala Leu Ile		
1	5	10
ggc agc ctg atg ctg ctg ctc ctc ctg cgt atg ctc tgg tgc cca gcc	157	
Gly Ser Leu Met Leu Leu Leu Leu Arg Met Leu Trp Cys Pro Ala		
15	20	25
gac gcg cct gcc cgc tcc agg ctg ttg atg gag gga agc aga gag gac	205	
Asp Ala Pro Ala Arg Ser Arg Leu Leu Met Glu Gly Ser Arg Glu Asp		
30	35	40
acc agt ggt acc tca gct gca ctg aag aca ctc tgg agc cag aca acc	253	
Thr Ser Gly Thr Ser Ala Ala Leu Lys Thr Leu Trp Ser Pro Thr Thr		
45	50	55

ccg gta cca cgc acc agg aac agc aca tat ctg gat gag aag aca acc	301
Pro Val Pro Arg Thr Arg Asn Ser Thr Tyr Leu Asp Glu Lys Thr Thr	
60 65 70 75	
caa ata aca gag aaa tgc aaa gat ctg caa tat agc ttg aac tct tta	349
Gln Ile Thr Glu Lys Cys Lys Asp Leu Gln Tyr Ser Leu Asn Ser Leu	
80 85 90	
tct aac aaa acg aga cgg tac tct gag gat gac tac ctc cag acc atc	397
Ser Asn Lys Thr Arg Arg Tyr Ser Glu Asp Asp Tyr Leu Gln Thr Ile	
95 100 105	
aca aac ata cag aga tgc cca tgg aac cgg caa gca gaa gaa tat gac	445
Thr Asn Ile Gln Arg Cys Pro Trp Asn Arg Gln Ala Glu Glu Tyr Asp	
110 115 120	
aat ttt aga gca aaa ctg gct tcc tgt tgc gat gcc att caa gac ttc	493
Asn Phe Arg Ala Lys Leu Ala Ser Cys Cys Asp Ala Ile Gln Asp Phe	
125 130 135	
gtg gtt tcc cag aac aac act cca gtg ggg act aac atg agc tac gag	541
Val Val Ser Gln Asn Asn Thr Pro Val Gly Thr Asn Met Ser Tyr Glu	
140 145 150 155	
glg gaa agc aag aaa cac atc ccc att cga gag aac att ttc cac atg	589
Val Glu Ser Lys Lys His Ile Pro Ile Arg Glu Asn Ile Phe His Met	
160 165 170	
ttt cca gtg tgc cag cct ttt gtg gac tat ccc tat aac cag tgt gca	637
Phe Pro Val Ser Gln Pro Phe Val Asp Tyr Pro Tyr Asn Gln Cys Ala	
175 180 185	
glg gtt ggt aat ggg gga att ctc aac aag tct ctc tgc gga gca gaa	685
Val Val Gly Asn Gly Gly Ile Leu Asn Lys Ser Leu Cys Gly Ala Glu	
190 195 200	

att gat aaa tct gac ttc gtc ttc agg tgt aac ctc ccc cca atc aca	733
Ile Asp Lys Ser Asp Phe Val Phe Arg Cys Asn Leu Pro Pro Ile Thr	
205 210 215	
ggg agc gct agt aaa gat gtt gga agc aaa aca aat ctt gtg act gtc	781
Gly Ser Ala Ser Lys Asp Val Gly Ser Lys Thr Asn Leu Val Thr Val	
220 225 230 235	
aat ccc agc att ata acc ctg aag tac cag aat ttg aag gag aag aaa	829
Asn Pro Ser Ile Ile Thr Leu Lys Tyr Gln Asn Leu Lys Glu Lys Lys	
240 245 250	
gca cag ttt ttg gag gac atc tcc acc tat gga gat gca ttc ctc ctc	877
Ala Gln Phe Leu Glu Asp Ile Ser Thr Tyr Gly Asp Ala Phe Leu Leu	
255 260 265	
ctg cca gca ttt tcc tat cgg gcc aac aca ggc atc tct ttt aaa gtc	925
Leu Pro Ala Phe Ser Tyr Arg Ala Asn Thr Gly Ile Ser Phe Lys Val	
270 275 280	
tac caa aca ctc aaa gag tca aaa atg agg caa aag gtt ctc ttc ttc	973
Tyr Gln Thr Leu Lys Glu Ser Lys Met Arg Gln Lys Val Leu Phe Phe	
285 290 295	
cat ccc agg tac ctg aga cac ctc gct ctt ttc tgg aga act aaa ggg	1021
His Pro Arg Tyr Leu Arg His Leu Ala Leu Phe Trp Arg Thr Lys Gly	
300 305 310 315	
gtg act gca tac cgc ttg tcc aca ggc ttg atg att gca agt gtc gct	1069
Val Thr Ala Tyr Arg Leu Ser Thr Gly Leu Met Ile Ala Ser Val Ala	
320 325 330	
gtg gaa ctg tgt gaa aac gtg aag ctc tac gga ttc tgg cct ttc tct	1117
Val Glu Leu Cys Glu Asn Val Lys Leu Tyr Gly Phe Trp Pro Phe Ser	
335 340 345	

aag act atc gaa gac acc cca ctc agt cac cac tac tat gat aac atg 1165
Lys Thr Ile Glu Asp Thr Pro Leu Ser His His Tyr Tyr Asp Asn Met
350 355 360
tta cct aag cat ggt ttc cac cag atg cct aaa gaa tac agc caa atg 1213
Leu Pro Lys His Gly Phe His Gln Met Pro Lys Glu Tyr Ser Gln Met
365 370 375
ctc cag ctc cat atg aga gga atc ctc aaa ctg caa ttc agc aaa tgt 1261
Leu Gln Leu His Met Arg Gly Ile Leu Lys Leu Gln Phe Ser Lys Cys
370 385 390 395
gaa acg get taa cgtttct tagaaggaga ataatttcag gaggtggagi 1310
Glu Thr Ala
398
ggaatgtgtca cagcatctcc aaaaagccaa tagaagaagg cacagagaaa gcatgaatt 1370
caaaggcgct cccccacttg tctagaccac agccaccgc cccactcac ttgcagcct 1430
ccacagatca ctcattctca ccttcaacgt tctttctctg agaatagaga ccaaacatc 1490
agacttggat aagtaaatg agataatttt tcaaalcatc atagaatttg atttgagcca 1550
gggtctctca gaatgcttcc ttgttcttat ccatgatage catteccacc ttatcagag 1610
tggtaatgaa actgtgcaat tgtgccaaag accctttctg aagagaatgt ctgaatcatg 1670
cgccgagtgt ttacacacag ctcttctctt ataaataaat ccttcccatc ctcctccta 1730
glagagtaca gaaacaaat acccttgatg attcaggaag aaaagtcttt ttactttagc 1790
aatgtgccgt cttctgaltc agttcgcttg tgacattaaag ctgggttggg gttttggtg 1850
gatttggggc gtttcttcac ttcttttgc tatatcttcc ttacctttat cagtttgat 1910
tcgagcttcc tgccttggga ttctgcaatt ctctctccca ctgacaggat caactcaatg 1970
acataaagta gtccaacat ccatgcttcc tcacatgttt tatccataaa gttactcatc 2030
tgattttatt taaaatagtg aacatctact tgatatcaga ccgaggacc atctccatt 2090
ggagaatatg aagatatgtg cactggcaga aaagcaggtg tgtgccatta attgataaga 2150
taccacaage atcatcatgc cagttatgaa cacagtgtg aaagatcat agacaggggt 2210

ggtaaaatct gatcccgta gaataaaactt cagtgtagct atttcaggga agagttaatt 2270
 tcacaattaa aactagtaaa tgaaccaatt ctaggcaca ttaagtgat tctgagtaaa 2330
 agaaggaggaa cagcaggaga aagctgttcg cttggtctg attacccaaa tgagcatgct 2390
 ggaaggaggl tglgggcta cgtlaaaacc tctgcgtagg gagagagtae agtgcattag 2450
 tgtggcggtt ttgtccaca ctctgaagg gtgagtaatt cagagccaat cacaicaca 2510
 ggatggacac acctaatca tcacttcagg gggagatgaa tgcattcatg agaattaca 2570
 ctataagct aagcatcagt tttagtaaa atttagtag atgttaata tgaacatttt 2630
 ataccctta ctaatgccc acgacacct ttaattgaa gcacatttat ttaataagt 2690
 acttgacatt aaatgcttat gtctgtatat tctgtctat calcalitt cccaaaagt 2750
 aagagcatag gagatgaggc ctacatgcca agaaaactat aaattttact cttaattct 2810
 tacttgacc agcttgttgt ttaacaagt cttttttgaa gagacagcac ccgtgaatt 2870
 ctctattctg atacagtgc accttgtatt taacatttgt aatgttgttt caagtttaca 2930
 tctctttat ctttttatag caaatcaaac gtattagctt cagaaattta tcagaagtic 2990
 atatataat attttgcaa ggtlaaaagg cttttttgtt aaataaata aaatttata 3050
 tttttcttg atgaatagag gctcttttat gctgtgcta algaacclaa ttagctttaa 3110
 attatctct agcaacattg gtcagtttc aalcagcta ttagcaaaaa aaaaaa 3166

[0056]

<210> 3

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic DNA

<400> 3

ctttcttgga gaactaaagg 20

[0057]

<210> 4

<211> 20

<212> DNA

<213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic DNA
 <400> 4
 aattgcagtt tgaggattcc 20
 [0058]
 <210> 5
 <211> 20
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: Synthetic DNA
 <400> 5
 tggctcagga tgagatcggg 20
 [0059]
 <210> 6
 <211> 22
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: Synthetic DNA
 <400> 6
 tactagecgt ccctgtgatt gg 22
 [0060]
 <210> 7
 <211> 30
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: Synthetic DNA
 <400> 7
 tgetctcgag cccagccgac gcgccgcc 30
 [0061]
 <210> 8
 <211> 30
 <212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic DNA

<400> 8

tattctcgag ctaagaaacg ttaagcgtt 30

[Brief Description of the Drawings]

[Fig.1]

Figure 1 shows the nucleotide sequences of ST8Sia VI cDNA, and the deduced amino acid sequences. A transmembrane domain is underlined, sialyl motif L is double-underlined, and sialyl motif S is dashed-underlined. Histidine and glutamic acid, which are conserved in sialyl motif VS, are boxed. Asparagine residues of the potential N-linked glycosylation sites are overlined.

[Fig.2]

Figure 2 shows a comparison made among the amino acid sequences of mouse sialyltransferases ST8Sia I, ST8Sia V, and ST8Sia VI. The conserved amino acid residues among these sialyltransferases are boxed. Sialyl motif L is double-underlined, and sialyl motif S is dashed-underlined. The conserved histidine and glutamic acid residues in sialyl motif VS are marked with asterisks.

[Fig.3]

Figure 3 shows an analysis of linkage of sialic acid which was introduced into GM3 by PA-ST8Sia VI, and a TLC immunostaining of the reaction product.

A, GM3 was sialylated with [¹⁴C]-NeuAc by PA-ST8Sia VI, and was treated with α 2,3-, and α 2,6-linkage specific sialidase (NANase II) or with α 2,3-, α 2,6-, α 2,8-, and α 2,9-linkage specific sialidase (NANase III), and then the reaction products were analyzed by HPTLC. The obtained results are shown.

B, GM3 was sialylated by PA-ST8Sia VI, and the reaction product was analyzed by TLC immunostaining. The obtained results are shown. Lane 1, GD3 (1 μ g); lane 2, GM3 (1 μ g); and lane 3, the reaction product. The reaction product was reacted with an


anti-GD3 monoclonal antibody KM641 and peroxidase-conjugated anti-mouse IgG + IgM (H+L) antibody, and then detected using an ECL system.

[Fig.4]


In Figure 4, Fetuin was [^{14}C]-NeuAc-incorporated by ST8Sia III or ST8Sia VI and then treated with *N*-glycanase.

The [^{14}C]-NeuAc-incorporated Fetuin was treated with *N*-glycanase, and the treated product was analyzed by SDS-PAGE. Thereafter, it was visualized with a BAS2000 radio image analyzer.

[Fig.3]



[Fig.4]



[Name of Document] ABSTRACT

[Abstract]

[Object] To provides a novel O-glycan α 2,8-sialyltransferase showing high activity toward O-glycans.

[Means for Solution] O-glycan α 2,8-sialyltransferase, which is characterized in that it has the following substrate specificity and substrate selectivity.

Substrate specificity: the substrates of the enzyme are glycoconjugates having a Sia α 2,3(6)Gal structure (wherein Sia represents sialic acid and Gal represents galactose) at the terminus thereof.

Substrate selectivity: the enzyme incorporates sialic acids into O-glycans more preferentially than into glycolipids or N-glycans.

[Selected Drawing] None